arXiv:hep-th/9605123v1 17 May 1996

RIMS-972
April 1994

Quantum Knizhnik–Zamolodchikov equation for $U_q(\widehat{\mathfrak{sl}_n})$ and integral formula

Quantum Knizhnik–Zamolodchikov equation
for $U_q(\widehat{\mathfrak{sl}_n})$ and integral formulaTakeo Kojima[1] and Yas-Hiro Quano[1,2]

[1] *Research Institute for Mathematical Sciences, Kyoto University, Kyoto 606, Japan*

[2] *Institute for Nuclear Study, University of Tokyo, Tanashi, Tokyo 188, Japan*

Abstract

Presented is an integral formula for solutions to the quantum Knizhnik–Zamolodchikov equation of level 0 associated with the vector representation of $U_q(\widehat{\mathfrak{sl}_n})$. This formula gives a generalization of both our previous work for $U_q(\widehat{\mathfrak{sl}_2})$ and Smirnov's formula for form factors of $SU(n)$ chiral Gross-Neveu model.

1 Introduction

In our previous paper [1] we gave an integral formula for solutions to the quantum Knizhnik–Zamolodchikov (KZ) equation [2] for the quantum affine algebra $U_q(\widehat{\mathfrak{sl}}_2)$ when the spin is 1/2, the level is 0 and $|q| < 1$. The present paper is a $U_q(\widehat{\mathfrak{sl}}_n)$ generalization of [1]. Our idea is based on Ref.[3]. Instead of solving the quantum KZ equation we consider a system of difference equations for a vector-valued function in N variables $(z_1, \cdots, z_N)$ which takes values in the N-fold tensor product of the vector representation $V = \mathbb{C}^n$ of $U_q(\widehat{\mathfrak{sl}}_n)$.

For a fixed complex number q satisfying $0 < |q| < 1$, let $R(z) \in \mathrm{End}(V \otimes V)$ be the standard trigonometric R-matrix associated with the vector representation $V \cong \mathbb{C}v_1 \oplus \cdots \oplus \mathbb{C}v_n$ of $U_q(\widehat{\mathfrak{sl}}_n)$. The matrix $R(z)$ satisfying the Yang-Baxter equation and the unitarity relation is given as follows:

$$R(z)v_{\varepsilon_1'} \otimes v_{\varepsilon_2'} = \sum_{\varepsilon_1,\varepsilon_2} v_{\varepsilon_1} \otimes v_{\varepsilon_2} R^{\varepsilon_1\varepsilon_2}_{\varepsilon_1'\varepsilon_2'}(z),$$

where the nonzero entries are

$$\begin{aligned}
R^{\varepsilon\varepsilon}_{\varepsilon\varepsilon}(z) &= 1, \\
R^{\varepsilon\varepsilon'}_{\varepsilon\varepsilon'}(z) &= b(z) = \frac{(1-z)q}{1-zq^2}, \quad \text{if } \varepsilon \neq \varepsilon', \\
R^{\varepsilon\varepsilon'}_{\varepsilon'\varepsilon}(z) &= \begin{cases} \dfrac{(1-q^2)z}{1-zq^2} & \text{if } \varepsilon < \varepsilon', \\ \dfrac{(1-q^2)}{1-zq^2} & \text{if } \varepsilon > \varepsilon'. \end{cases}
\end{aligned}$$

In statistical mechanics language each entry of the R-matrix is a local Boltzmann weight for a single vertex with bond states $i, j, k, l \in \mathbb{Z}_n$:

$R^{ik}_{jl}(z_1/z_2) =$ 

,

where each line carries a spectral parameter.

In what follows we shall work with the tensor product of finitely many V's. Following the usual convention we let $R_{jk}(z)$ $(j \neq k)$ signify the operator on $V^{\otimes N}$ acting as $R(z)$ on the (j,k)-th tensor components and as identity on the other components. In particular we have $R_{kj}(z) = P_{jk}R_{jk}(z)P_{jk}$, where $P \in \mathrm{End}(V \otimes V)$ stands for the transposition $P(x \otimes y) = y \otimes x$.

The equations we are concerned with in this paper are (1) R-matrix symmetry and (2) deformed cyclicity, for a function $G(z_1, \cdots, z_N) \in V^{\otimes N}$:

$$(1) \qquad P_{j\,j+1}G(\cdots, z_{j+1}, z_j, \cdots) = R_{j\,j+1}(z_j/z_{j+1})G(\cdots, z_j, z_{j+1}, \cdots), \tag{1.1}$$

$$(2) \qquad P_{12}\cdots P_{N-1N}G(z_2,\cdots,z_N,z_1q^{-2n}) = D_1G(z_1,\cdots,z_N). \tag{1.2}$$

In (1.2) D_1 is an operator acting on the first component as $D = \mathrm{diag}(\delta_1,\cdots,\delta_n)$, whose entries will be specified below, and as identity on the other ones. These are two of the axioms that form factors in integrable models should satisfy [4]. Smirnov [4] also pointed out that (1.1) and (1.2) imply the quantum KZ equation of level 0 [2]

$$\begin{aligned} G(z_1,\cdots,z_jq^{2n},\cdots,z_N) = R_{j-1\,j}(z_{j-1}/z_jq^{2n})^{-1}\cdots R_{1\,j}(z_1/z_jq^{2n})^{-1}D_j^{-1} \\ \times R_{j\,N}(z_j/z_N)\cdots R_{j\,j+1}(z_j/z_{j+1})G(z_1,\cdots,z_j,\cdots,z_N). \end{aligned} \tag{1.3}$$

Throughout this article the functions we consider are not necessarily single valued in z_j but are meromorphic in the variable $\log z_j$. Accordingly the shift $z_j \to z_jq^{-2n}$ as in (1.2) is understood to mean $\log z_j \to \log z_j - 2n\log q$.

In the sequel we set $\tau = q^{-1}$. Define the components of G by

$$G(z_1,\cdots,z_N) = \sum_{\varepsilon_j=1}^{n} v_{\varepsilon_1}\otimes\cdots\otimes v_{\varepsilon_N}G^{\varepsilon_1\cdots\varepsilon_N}(z_1,\cdots,z_N). \tag{1.4}$$

Then the equation (1.1) reads as

$$G^{\cdots\varepsilon\varepsilon\cdots}(\cdots,z_j,z_{j+1},\cdots) = G^{\cdots\varepsilon\varepsilon\cdots}(\cdots,z_{j+1},z_j,\cdots), \tag{1.5}$$

$$\begin{aligned} G^{\cdots\varepsilon\varepsilon'\cdots}(\cdots,z_j,z_{j+1},\cdots) &= \frac{z_j - z_{j+1}\tau^2}{(z_j-z_{j+1})\tau}G^{\cdots\varepsilon'\varepsilon\cdots}(\cdots,z_{j+1},z_j,\cdots) \\ &- \frac{(1-\tau^2)z_{jj+1}^{\varepsilon\varepsilon'}}{(z_j-z_{j+1})\tau}G^{\cdots\varepsilon'\varepsilon\cdots}(\cdots,z_j,z_{j+1},\cdots), \end{aligned} \tag{1.6}$$

where

$$z_{jj+1}^{\varepsilon\varepsilon'} = \begin{cases} z_j & \text{if } \varepsilon < \varepsilon', \\ z_{j+1} & \text{if } \varepsilon > \varepsilon', \end{cases}$$

and (1.2) reads as

$$G^{\varepsilon_2\cdots\varepsilon_N\varepsilon_1}(z_2,\cdots,z_N,z_1\tau^{2n}) = \delta_{\varepsilon_1}\ G^{\varepsilon_1\varepsilon_2\cdots\varepsilon_N}(z_1,z_2,\cdots,z_N). \tag{1.7}$$

Note that the singularity at $z_j = z_{j+1}$ in (1.6) is spurious. The equations (1.5–1.7) split into blocks, each involving components such that

$$\sharp\{j|\varepsilon_j = i\} = m_i, \quad N = \sum_{i=1}^{n} m_i.$$

In the present paper we restrict ourselves to the case $m_1 = \cdots = m_n = m$ and hence $N = mn$. According to this restriction, we set $\delta_i = \tau^{m(1-n)+2(1-i)}$.

We use the abbreviation $z^{(j)} = (z_1^{(j)}, \cdots, z_m^{(j)})$. Consider the extreme component

$$G^{\overbrace{n\cdots n}^{m}\cdots\overbrace{1\cdots 1}^{m}}(z^{(n)}|\cdots|z_1^{(1)}) = H(z^{(n)}|\cdots|z^{(1)}). \tag{1.8}$$

Because of (1.5) this function is symmetric in the variables $z^{(1)}$'s, $\cdots$, $z^{(n)}$'s, separately. The equation (1.6) tells that all the components with fixed m are uniquely determined from H. Conversely given any such H the Yang-Baxter equation guarantees that (1.1) can be solved consistently under the condition (1.8).

We wish to find an integral formula of the form

$$H(z_1, \cdots, z_N) = (S_{MN}F)(z_1, \cdots, z_N), \tag{1.9}$$

where S_{MN} stands for the following integral transform

$$(S_{MN}F)(z_1, \cdots, z_N) = \oint_C dx_1 \cdots \oint_C dx_M F(x_1, \cdots, x_M|z_1, \cdots, z_N)\Psi(x_1, \cdots, x_M|z_1, \cdots, z_N). \tag{1.10}$$

The notation is explained below.

The kernel Ψ has the form

$$\Psi(x_1, \cdots, x_M|z_1, \cdots, z_N) = \vartheta(x_1, \cdots, x_M|z_1, \cdots, z_N) \prod_{\mu=1}^{M}\prod_{j=1}^{N} \psi\Big(\frac{x_\mu}{z_j}\Big), \tag{1.11}$$

where

$$\psi(z) = \frac{1}{(zq^{n-1}; q^{2n})_\infty (z^{-1}q^{n-1}; q^{2n})_\infty}, \qquad (z;p)_\infty = \prod_{k=0}^{\infty}(1 - zp^k). \tag{1.12}$$

Assume that the function ϑ is anti-symmetric and holomorhpic in the $x_\mu \in \mathbb{C}\backslash\{0\}$, and is symmetric and meromorphic in the $\log z_j \in \mathbb{C}$, possessing the following transformation property

$$\begin{aligned} \vartheta(x_1, \cdots, x_M|z_1, \cdots, z_j\tau^{2n}, \cdots, z_N) &= \vartheta(x_1, \cdots, x_M|z_1, \cdots, z_N) \prod_{\mu=1}^{M} \frac{-z_j\tau^{n-1}}{x_\mu}, \\ \vartheta(x_1, \cdots, x_\mu\tau^{2n}, \cdots, x_M|z_1, \cdots, z_N) &= \vartheta(x_1, \cdots, x_M|z_1, \cdots, z_N) \prod_{j=1}^{N} \frac{-x_\mu\tau^{n-1}}{z_j}. \end{aligned} \tag{1.13}$$

The function ϑ is otherwise arbitrary, and the choice of ϑ's corresponds to that of solutions. The integration $\oint_C dx_\mu$ is along a simple closed curve $C = C(z_1, \cdots, z_N)$ oriented anti-clockwise, which encircles the points $z_j\tau^{-n+1-2nk}(1 \le j \le N, k \ge 0)$ but not $z_j\tau^{n-1+2nk}(1 \le j \le N, k \ge 0)$. Finally

$$F(x_1, \cdots, x_M|z^{(n)}|\cdots|z^{(1)}) = \frac{\Delta^{(m)}(x_1, \cdots, x_M|z^{(n)}|\cdots|z^{(1)})}{\prod_{\substack{k,k'=1\\k<k'}}^{n}\prod_{j=1}^{m}\prod_{j'=1}^{m}(z_j^{(k)} - z_{j'}^{(k')}\tau^2)}, \tag{1.14}$$

where $\Delta^{(m)}$ is a certain homogeneous polynomial to be determined, antisymmetric in the variables $(x_1, \cdots, x_n)$ and symmetric in the variables $z^{(1)}$'s, $\cdots$, $z^{(n)}$'s, separately.

The rest of the paper is organized as follows. In Section 2 we introduce a special basis of V in terms of the quantum monodromy operators. In Section 3 we describe the main theorem of the present paper. The subsequent two sections are devoted to proof of it. Section 4 is for $m = 1$ and Section 5 is for general case. In section 6 we discuss the relation among other works and ours.

2 Quantum monodromy operators and the special basis

We shall construct a special basis $\{w_{\alpha_1 \cdots \alpha_N}(z_1, \cdots, z_N)\}$ with $\alpha_j \in \{1, 2, \cdots, n\}$ of $V^{\otimes N}$ depending on the parameter $(z_1, \cdots, z_N)$, which satisfy

$$P_{jj+1} w_{\cdots \alpha_{j+1} \alpha_j \cdots}(\cdots, z_{j+1}, z_j, \cdots) = R_{jj+1}(z_j/z_{j+1}) w_{\cdots \alpha_j \alpha_{j+1} \cdots}(\cdots, z_j, z_{j+1}, \cdots). \tag{2.1}$$

The procedure goes as follows. Define the quantum monodromy operator $\mathcal{T}_{\varepsilon\varepsilon'}(z_1, \cdots, z_N|t) \in \mathrm{End}(V^{\otimes N})$ by

$$R_{1\,N+1}(z_1/t) \cdots R_{N\,N+1}(z_N/t) = (\mathcal{T}_{\varepsilon\varepsilon'}(z_1, \cdots, z_N|t))_{1 \le \varepsilon, \varepsilon' \le n}\,. \tag{2.2}$$

Here the $n \times n$ matrix structure is defined relative to the base $v_1, \cdots, v_n$ of the $(N+1)$-th tensor component of $V^{\otimes (N+1)}$:

$\mathcal{T}_{\varepsilon\varepsilon'}(z_1, \cdots, z_N|t) =$



For $\alpha = (\alpha_1, \cdots, \alpha_N)$ with $\alpha_j \in \{1, \cdots, n\}$, set $J_i^\alpha = \{\ j \mid \alpha_j = i\}$ and $\{z_1^{(i)}, \cdots, z_m^{(i)}\} = \{z_j | \alpha_j = i\}$. where $1 \le i \le n$, and set

$$\begin{aligned} w_\alpha(z_1, \cdots, z_N) &= \prod_{l=1}^{m} \mathcal{T}_{1n}(z_1, \cdots, z_N | z_l^{(n)}) \cdots \prod_{l=1}^{m} \mathcal{T}_{12}(z_1, \cdots, z_N | z_l^{(2)}) \Omega, \\ \Omega &= v_1 \otimes \cdots \otimes v_1 \quad \in V^{\otimes N}. \end{aligned} \tag{2.3}$$

Then $w_\alpha(z_1, \cdots, z_N)$ is visualized as follows:

$$w_\alpha(z_1,\cdots,z_N)=\sum v_{i_1}\otimes\cdots\otimes v_{i_N}$$



Let us introduce the ordered indices set $(\alpha_1,\cdots,\alpha_N)>(\varepsilon_1,\cdots,\varepsilon_N)$ if and only if $\alpha_i=\varepsilon_i,(1\le i\le k)$ and $\alpha_{k+1}>\varepsilon_{k+1}$. Define the components of the basis by

$$w_{\alpha_1\cdots\alpha_N}(z_1,\cdots,z_N)=\sum_{\varepsilon_1,\cdots,\varepsilon_N} v_{\varepsilon_1}\otimes\cdots\otimes v_{\varepsilon_N} w^{\varepsilon_1\cdots\varepsilon_N}_{\alpha_1\cdots\alpha_N}(z_1,\cdots,z_N), \tag{2.4}$$

then we have

$$w^{\varepsilon_1\cdots\varepsilon_N}_{\alpha_1\cdots\alpha_N}(z_1,\cdots,z_N)=0,\quad \text{if}\quad (\alpha_1,\cdots,\alpha_N)<(\varepsilon_1,\cdots,\varepsilon_N). \tag{2.5}$$

Furthermore for $\beta=(n,\cdots,n,\cdots,2,\cdots,2,1,\cdots,1)$ we have

$$w^\beta_\beta(z_1,\cdots,z_N)=\prod_{\substack{k\in J^\beta_i,l\in J^\beta_j\\ i<j}} b(z_k/z_l). \tag{2.6}$$

From (2.1), (2.5) and (2.6) we obtain the following explicit formula

$$G(z_1,\cdots,z_N)=\sum_\alpha w_\alpha(z_1,\cdots,z_N)H(\{z_j\}_{j\in J^\alpha_n}|\cdots|\{z_j\}_{j\in J^\alpha_1})\prod_{\alpha_i<\alpha_j}\frac{1}{b(z_i/z_j)}.$$

For $2\le p\le n$ and $n\ge i_1>\cdots>i_p\ge 1$ set

$$v^{(i_1\cdots i_p)}=\sum_{\sigma\in\mathfrak{S}_p}(-\tau)^{l(\sigma)}v_{\sigma(i_1)}\otimes\cdots\otimes v_{\sigma(i_p)},$$

where $l(\sigma)$ is the minimum number of permutations such that

$$v_{\sigma(i_1)} \otimes \cdots \otimes v_{\sigma(i_p)} = \Big(\prod P_{jj+1}\Big)\, v_{i_1} \otimes \cdots \otimes v_{i_p}.$$

For example

$$\begin{aligned} v^{(21)} &= v_2 \otimes v_1 - \tau v_1 \otimes v_2, \\ v^{(321)} &= v_3 \otimes v_2 \otimes v_1 - \tau(v_3 \otimes v_1 \otimes v_2 + v_2 \otimes v_3 \otimes v_1) \\ &+ \tau^2(v_1 \otimes v_3 \otimes v_2 + v_2 \otimes v_1 \otimes v_3) - \tau^3 v_1 \otimes v_2 \otimes v_3. \end{aligned}$$

For $i \in \{i_1, \cdots, i_p\}$ one can easily check the following formulae

$$R_{1p+1}(z)R_{2p+1}(z\tau^2)\cdots R_{pp+1}(z\tau^{2p-2})v^{(i_1\cdots i_p)} \otimes v_i = \prod_{j=1}^{p-1} b(z\tau^{2j})v^{(i_1\cdots i_p)} \otimes v_i. \tag{2.7}$$

The poles of $w_\alpha(z)$ exist only at $z_j = z_i\tau^2$ for $i < j$ and $\alpha_i > \alpha_j$. Thus we have

$$\begin{aligned} &\mathrm{Res}_{z_2=z_1\tau^2}\cdots\mathrm{Res}_{z_n=z_{n-1}\tau^2} w_{\alpha_1\cdots\alpha_n}(z_1,\cdots,z_n) \\ =\ & \delta_{\alpha_1 n}\delta_{\alpha_2 n-1}\cdots\delta_{\alpha_n 1} z_1^{n-1}(\tau^2-1)^{n-1}\tau^{(n-1)^2}\,[n-1]!\ \ v^{(n\cdots 21)}, \end{aligned} \tag{2.8}$$

where

$$[k]! = [k]\cdots[1]\,, \qquad [k] = \frac{\tau^k - \tau^{-k}}{\tau - \tau^{-1}}.$$

Furthermore we obtain the recursive residue formula

$$\begin{aligned} &\mathrm{Res}_{z_2=z_1\tau^2}\cdots\mathrm{Res}_{z_n=z_{n-1}\tau^2} w_{\alpha_1\cdots\alpha_N}(z_1,\cdots,z_N) = \textstyle\prod_{\substack{\alpha_i>\alpha_j \\ i\le n<j}} b(z_j/z_i) \\ \times\ & \Big(\mathrm{Res}_{z_2=z_1\tau^2}\cdots\mathrm{Res}_{z_n=z_{n-1}\tau^2} w_{n\cdots 1}(z_1,\cdots,z_n)\Big) \otimes w_{\alpha_{n+1}\cdots\alpha_N}(z_{n+1},\cdots,z_N), \end{aligned} \tag{2.9}$$

for $\alpha_i = n+1-i$ $(1 \le i \le n)$. By combining (2.1) and (2.9) and using (2.7) we have the useful expression

$$\begin{aligned} &\mathrm{Res}_{z_m^{(n-1)}=z_m^{(n)}\tau^2}\cdots\mathrm{Res}_{z_m^{(1)}=z_m^{(2)}\tau^2} w_\alpha^{\overbrace{n\cdots n}^{m}\cdots\hat{i}\cdots\hat{i}\cdots\overbrace{1\cdots 1}^{m}\overbrace{i\cdots i}^{m}}(z^{(n)}|\cdots|z^{(1)}) \\ =\ & \prod_{k=2}^{n}\prod_{l=1}^{m-1}\prod_{j=n+1-k}^{n-1} b(z_m^{(n)}\tau^{2j}/z_l^{(k)}) \prod_{\substack{\alpha_{mk+l}<\alpha_{jm} \\ 1\le l<m}} b(z_l^{(n-k)}/z_m^{(j)}) \\ \times\ & (z_m^{(n)})^{n-1}(-\tau)^{i-1}(\tau^2-1)^{n-1}\tau^{(n-1)^2}[n-1]! \\ \times\ & v^{(n\cdots 21)} \otimes w_{\alpha'}^{\overbrace{n\cdots n}^{m-1}\cdots\hat{i}\cdots\hat{i}\cdots\overbrace{1\cdots 1}^{m-1}\overbrace{i\cdots i}^{m-1}}(z'^{(n)}|\cdots|z'^{(1)}). \end{aligned} \tag{2.10}$$

for $\alpha' = (\alpha_1, \cdots, \hat{\alpha_m}, \cdots, \alpha_{\hat{2m}}, \cdots, \alpha_{nm-1})$ and $\alpha_{mi} = n+1-i$ $(1 \le i \le n)$. Here we use the abbreviation $z'^{(i)} = (z_1^{(i)}, \cdots, z_{m-1}^{(i)})$.

3 Main theorem

Now we present the main theorem of the present paper. In what follows we use the abbreviation

$$\begin{aligned} z^{(j)} &= (z_1^{(j)}, \cdots, z_{m-1}^{(j)}, z_m^{(j)}) = (z'^{(j)}, z_m^{(j)}), \\ z^{(j)}\tau^{\pm 1} &= (z_1^{(j)}\tau^{\pm 1}, \cdots, z_{m-1}^{(j)}\tau^{\pm 1}, z_m^{(j)}\tau^{\pm 1}) = (z'^{(j)}\tau^{\pm 1}, z_m^{(j)}\tau^{\pm 1}), \\ x &= (x_1, \cdots, x_M). \end{aligned}$$

The polynomial $\Delta^{(m)}$ in (1.14) is given by

$$\Delta^{(m)}(x|z^{(n)}|\cdots|z^{(2)}|z^{(1)}) = \det\left(A_\lambda^{(m)}(x_\mu|z^{(n)}|\cdots|z^{(2)}|z^{(1)})\right)_{1\le\lambda,\mu\le M} \tag{3.1}$$

where $M = M_m = (n-1)m - 1, N = mn$. The entries of the $M \times M$ matrix $A^{(m)}$ is defined as follows. Let us introduce the polynomial

$$f_\lambda^{(N)}(y|z_1, \cdots, z_N) = \sum_{\kappa=0}^{\lambda-1}(-1)^\kappa((y\tau)^{\lambda-\kappa} - (y\tau^{-1})^{\lambda-\kappa})\sigma_\kappa(z_1, \cdots, z_N),$$

where $\sigma_\kappa(z_1, \cdots, z_n)$ denotes the κ-th elementary symmetric polynomials:

$$\prod_{j=1}^n (t + z_j) = \sum_{\kappa=0}^n \sigma_\kappa(z_1, \cdots, z_n)t^{n-\kappa}.$$

Note that for $\alpha > 0$

$$f_{N+\alpha}^{(N)}(y|z_1, \cdots, z_N) = y^\alpha \left\{ \tau^\alpha \prod_{j=1}^N (y\tau - z_j) - \tau^{-\alpha} \prod_{j=1}^N (y\tau^{-1} - z_j) \right\}. \tag{3.2}$$

Define the polynomial

$$\begin{aligned} &A_\lambda^{(m)}(x|z^{(n)}|\cdots|z^{(1)}) = \sum_{k=1}^n \prod_{j=1}^m (x - z_j^{(k)}\tau^{2k-n-1})\times \\ &\times \frac{1}{x} f_\lambda^{((n-1)m)}(x\tau^{n+1-2k}|z^{(n)}\tau, \cdots, z^{(k+1)}\tau \overset{k}{\hat{}} z^{(k-1)}\tau^{-1}, \cdots, z^{(1)}\tau^{-1}). \end{aligned} \tag{3.3}$$

This is a homogeneous polynomial of degree $m + \lambda - 1$, symmetric with respect to $z^{(k)}$'s for each k, separately. By the construction (3.1) and (3.3), $\Delta^{(m)}$ is a homogeneous polynomial of degree $M_m m + M_m(M_m + 1)/2$ with correct symmetries. For $n = 3$, it reads as

$$\begin{aligned} A_\lambda^{(m)}(x|z^{(3)}|z^{(2)}|z^{(1)}) &= \prod_{j=1}^m (x - z_j^{(3)}\tau^2)\frac{1}{x} f_\lambda^{(2m)}(x\tau^{-2}|z^{(2)}\tau^{-1}, z^{(1)}\tau^{-1}) \\ &+ \prod_{j=1}^m (x - z_j^{(2)})\frac{1}{x} f_\lambda^{(2m)}(x|z^{(3)}\tau, z^{(1)}\tau^{-1}) \\ &+ \prod_{j=1}^m (x - z_j^{(1)}\tau^{-2})\frac{1}{x} f_\lambda^{(2m)}(x\tau^2|z^{(3)}\tau, z^{(2)}\tau). \end{aligned}$$

The following is the main theorem of this paper.

Theorem 3.1 *The integral formula*

$$G(z_1,\cdots,z_N)=\sum_{\alpha} w_\alpha(z_1,\cdots,z_N)H(\{z_j\}_{j\in J_n^\alpha}|\cdots|\{z_j\}_{j\in J_1^\alpha})\prod_{\alpha_i<\alpha_j}\frac{1}{b(z_i/z_j)}$$

satisfies (1.1) and (1.2) with $(\delta_1,\delta_2,\cdots,\delta_n)=(\tau^{-(n-1)m},\tau^{-(n-1)m-2},\cdots,\tau^{-(n-1)(m+2)})$*, where* H *is defined by (1.9–1.14) and* $w_\alpha(z_1,\cdots,z_N)$ *is defined by (2.2–2.3), respectively.*

First of all, we would like to notice the following lemmas:

Lemma 3.2 *The following holds:*

$$(S_{MN}\Delta^{(m)})(z^{(n)}|\cdots|z^{(2)}|z'^{(1)},z_0\tau^{2n})=(S_{MN}\tilde{\Delta}^{(m)})(x|z^{(n)}|\cdots|z^{(2)}|z'^{(1)},z_0),$$

where

$$\tilde{\Delta}^{(m)}(x|z^{(n)}|\cdots|z^{(2)}|z'^{(1)},z_0)=\det\left(\tilde{A_\lambda}^{(m)}(x_\mu|z^{(n)}|\cdots|z^{(2)}|z'^{(1)},z_0)\right)_{1\le\lambda,\mu\le n-2},$$

and

$$\begin{aligned}&\tilde{A}_\lambda^{(m)}(x|z^{(n)}|\cdots|z^{(2)}|z'^{(1)},z_0)=\frac{x-z_0\tau^{n-1}}{x-z_0\tau^{n+1}}A_\lambda^{(m)}(x|z^{(n)}|\cdots|z^{(2)}|z'^{(1)},z_0\tau^{2n})\\ +\ &\frac{z_0\tau^{n+1}}{x}\left\{\tau^{-2}\prod_{j=1}^{N-1}\frac{x\tau^{n-1}-z_j}{z_0\tau^2-z_j}-\frac{x-z_0\tau^{n-1}}{x-z_0\tau^{n+1}}\prod_{j=1}^{N-1}\frac{x\tau^{-n+1}-z_j}{z_0\tau^2-z_j}\right\}\\ \times\ &A_\lambda^{(m)}(z_0\tau^{n+1}|z^{(n)}|\cdots|z^{(2)}|z'^{(1)},z_0\tau^{2n}).\end{aligned}$$

It can be proved in a similar manner as in the case $n=2$, see [1].

To prove Theorem 3.1 it is enough to show (1.7) for n cases; i.e.,

$$\begin{aligned}&G^{\overbrace{n\cdots n}^{m}\cdots\hat{i}\cdots\hat{i}\cdots\overbrace{1\cdots 1}^{m}\overbrace{i\cdots i}^{m}}(z^{(n)}|\cdots|z'^{(1)},z_0\tau^{2n})\\ =\ &\delta_i\ G^{i\overbrace{n\cdots n}^{m}\cdots\hat{i}\cdots\hat{i}\cdots\overbrace{1\cdots 1}^{m}\overbrace{i\cdots i}^{m-1}}(z_0|z^{(n)}|\cdots|z'^{(1)}),\end{aligned}\tag{3.4}$$

for $i=1,\cdots,n$.

Let $z_j^{(k)}=z_{(n-k)m+j}$ for $n\ge k\ge 2$ and $z_j'^{(1)}=z_{(n-1)m+j}$. Let us define $J_i(j_1,\cdots,j_{i-1})$ recursively as follows:

$$\begin{aligned}J_1&=\{0,1,\cdots,m\}\ni j_1,\\ J_2(j_1)&=\{j_1,m+1,\cdots,2m\}\ni j_2,\\ &\vdots\\ J_{n-2}(j_1,\cdots,j_{n-3})&=\{j_{n-3},(n-3)m+1,\cdots,(n-2)m\}\ni j_{n-2}\\ J_{n-1}(j_1,\cdots,j_{n-2})&=\{j_{n-2},(n-2)m+1,\cdots,(n-1)m\}\ni j_{n-1}\end{aligned}\tag{3.5}$$

Set

$$\varphi^{(m)}(j_1,\cdots,j_{n-1}) = \prod_{k_1\in J_1\setminus\{j_1\}} \frac{z_{k_1}-z_0\tau^2}{(z_{j_1}-z_{k_1})\tau} \prod_{k_2\in J_2\setminus\{j_2\}} \frac{z_{k_2}-z_{j_1}\tau^2}{(z_{j_2}-z_{k_2})\tau}\cdots$$
$$\times \quad \cdots \prod_{k_{n-1}\in J_{n-1}\setminus\{j_{n-1}\}} \frac{z_{k_{n-1}}-z_{j_{n-2}}\tau^2}{(z_{j_{n-1}}-z_{k_{n-1}})\tau} \prod_{j=1}^{m-1} \frac{z_j'^{(1)}-z_{j_{n-1}}\tau^2}{(z_j'^{(1)}-z_0)\tau^2}.$$

The equation (3.4) for $i=1$ and $\delta_1=\tau^{m(1-n)}$ is satisfied if the following holds:

Proposition 3.3

$$\begin{aligned} &\sum_J \varphi^{(m)}(j_1,\cdots,j_{n-1})\Delta^{(m)}(x|\overset{j_1}{\hat{z_{J_1}}}|\cdots|\overset{j_{n-1}}{\hat{z_{J_{n-1}}}}|z_{j_{n-1}},z'^{(1)}) \\ = \quad &(-\tau)^{m(1-n)}\prod_{k=2}^{n}\prod_{j=1}^{m}\frac{z_j^{(k)}-z_0\tau^2}{z_j^{(k)}-z_0\tau^{2n-2}}\tilde{\Delta}(x|z^{(n)}|\cdots|z^{(2)}|z'^{(1)},z_0). \end{aligned} \tag{3.6}$$

In section 4 we prove Theorem 3.1 for $m=1$. In section 5 we verify Proposition 3.3 and also show that (3.4) for $i\neq 1$ reduces to Proposition 3.3, which imply Theorem 3.1 for any m.

4 The $m=1$ case

In this section we prove Theorem 3.1 for $m=1$. Notice that for $m=1$ $A_\lambda^{(m)}(x|z)$ coincides with

$$A_\lambda^{(1)}(x|z_1,\cdots,z_n) = \sum_{\kappa=0}^{\lambda}(-1)^\kappa x^{\lambda-\kappa}(\tau^{n(\lambda-\kappa)+\kappa}-\tau^{-n(\lambda-\kappa)-\kappa})\sigma_\kappa(z_1,\cdots,z_n). \tag{4.1}$$

and that it is linear with respect to z's. In this case $z_i = z_1^{(n+1-i)}$. In this section we use the abbreviations $\Delta^{(1)}=\Delta, \tilde{\Delta}^{(1)}=\tilde{\Delta}, A_\lambda^{(1)}=A_\lambda$. Since the polynomial $A_\lambda(x|z_1,\cdots,z_n)$ is symmetric with respect to the variable $(z_1,\cdots,z_n)$, by using (1.6) we obtain

$$\begin{aligned} G(z_1,\cdots,z_n)^{n\cdots\hat{i}\cdots 1i} &= (-\tau)^{1-i}H(z_1,\cdots,z_n), \\ G(z_1,\cdots,z_n)^{in\cdots\hat{i}\cdots 1} &= (-\tau)^{i-1}H(z_1,\cdots,z_n). \end{aligned} \tag{4.2}$$

Thus

$$G(z_2,\cdots,z_n,z_1\tau^{2n})^{n\cdots 21} = \delta_1 G(z_1,\cdots,z_n)^{1n\cdots 2}, \quad \delta_1=\tau^{1-n}, \tag{4.3}$$

implies

$$G(z_2,\cdots,z_n,z_1\tau^{2n})^{n\cdots\hat{i}\cdots 1i} = \delta_i G(z_1,\cdots,z_n)^{in\cdots\hat{i}\cdots 1}, \quad \delta_i=\delta_1\tau^{2-2i}. \tag{4.4}$$

Consequently, to prove Theorem 3.1 for $m=1$, it is enough to show (4.3).

Now we prepare the following lemmas.

Lemma 4.1

$$\tilde{\Delta}(x_1,\cdots,x_{n-2}|z_1,\cdots,z_n)|_{z_j=z_1\tau^{2n-2}}=0,\quad (j=2,3,\cdots,n). \tag{4.5}$$

Proof Note that

$$\begin{aligned}&\sum_{\lambda=0}^{n-3} z_1^{\lambda} A_{n-2-\lambda}(x|z_1\tau^{-1},z_1\tau,z_3,\cdots,z_n)\\ =\ &\frac{1}{x}\left\{(x\tau^{n-2}-z_1)\prod_{j=3}^{n}(x-z_j\tau^{-n+1})-(x\tau^{-n+2}-z_1)\prod_{j=3}^{n}(x-z_j\tau^{n-1})\right\}.\end{aligned}$$

Hence we obtain the linear dependence

$$\sum_{\lambda=0}^{n-3}(z_1\tau^{2n-1})^{\lambda}\tilde{A}_{n-2-\lambda}(x|z_1,z_1\tau^{2n-2},z_3,\cdots,z_n)=0,$$

which implies (4.5). □

Lemma 4.2

$$\begin{aligned}&\prod_{\mu=2}^{n-2}(x_\mu-z_1\tau)\Delta(z_1\tau^{-1},x_2,\cdots,x_{n-2}|z_1\tau^{-n},z_2,\cdots,z_n)\\ =\ &\prod_{\mu=2}^{n-2}(x_\mu-z_1\tau^{-1})\Delta(z_1\tau,x_2,\cdots,x_{n-2}|z_1\tau^{n},z_2,\cdots,z_n).\end{aligned} \tag{4.6}$$

Proof Let us first show that

$$\det(C_{\lambda,\mu})_{1\le,\mu,\lambda\le n-2}=\det(C'_{\lambda,\mu})_{1\le,\mu,\lambda\le n-2}, \tag{4.7}$$

where

$$C_{\lambda\mu}=\begin{cases}A_\lambda(z_1\tau^{-1}|z_1\tau^{-n},z_2,\cdots,z_n), & \text{if } \mu=1,\\ (x_\mu-z_1\tau)A_\lambda(x_\mu|z_1\tau^{-n},z_2,\cdots,z_n), & \text{if } \mu\neq 1,\end{cases}$$

$$C'_{\lambda\mu}=\begin{cases}A_\lambda(z_1\tau|z_1\tau^{n},z_2,\cdots,z_n), & \text{if } \mu=1,\\ (x_\mu-z_1\tau^{-1})A_\lambda(x_\mu|z_1\tau^{n},z_2,\cdots,z_n). & \text{if } \mu\neq 1.\end{cases}$$

Perform the following elementary transformations to the matrix $(C_{\lambda\mu})_{1\le\lambda,\mu\le n-2}$

$$\begin{aligned}&(1)(i\text{th row})-z_1\tau^{n-1}((i-1)\text{st row}),\quad (i=2,\cdots,n-2),\\ &(2)(i\text{th column})+z_1\tau((i-1)\text{st column}),\quad (i=2,\cdots,n-2),\end{aligned}$$

and to the matrix $(C'_{\lambda\mu})_{1\le\lambda,\mu\le n-2}$

$$
\begin{aligned}
&(1)(i\text{th row}) - z_1\tau^{-n+1}((i-1)\text{st row}), \quad (i=2,\cdots,n-2),\\
&(2)(i\text{th column}) + z_1\tau^{-1}((i-1)\text{st column}), \quad (i=2,\cdots,n-2).
\end{aligned}
$$

Then we have (4.7) and therefore (4.6). □

Because the polynomial $A_\lambda(z_1,\cdots,z_n)$ is symmetric with respect to $(z_1,\cdots,z_n)$, Proposition 3.3 holds if the following Proposition holds.

Proposition 4.3

$$
\tilde{\Delta}(x_1,\cdots,x_{n-2}|z_1,\cdots,z_n) = \prod_{j=2}^{n}\frac{z_j - z_1\tau^{2n-2}}{z_j - z_1\tau^2}\Delta(x_1,\cdots,x_{n-2}|z_1,\cdots,z_n).
$$

Proof Thanks to the linearity of the determinant we have

$$
\begin{aligned}
&\tilde{\Delta}(x_1,\cdots,x_{n-2}|z_1,\cdots,z_n)\\
=\;&\prod_{\mu=1}^{n-2}\frac{x_\mu - z_1\tau^{n-1}}{x_\mu - z_1\tau^{n+1}}\Delta(x_1,\cdots,x_{n-2}|z_1,\cdots,z_n)\\
+\;&\sum_{\nu=1}^{n-2}\prod_{\substack{\mu=1\\ \mu\neq\nu}}^{n-2}\frac{x_\mu - z_1\tau^{n-1}}{x_\mu - z_1\tau^{n+1}}g(x_\nu|z_1|z_2,\cdots,z_n)\\
\times\;&\Delta(x_1,\cdots,z_1\tau^{\overset{\nu}{\hat{n+1}}}\cdots x_{n-2}|z_1\tau^{2n},z_2,\cdots,z_n),
\end{aligned}
\tag{4.8}
$$

where

$$
g(x|z_1|z_2,\cdots,z_n) = \frac{z_1\tau^{n+1}}{x}\left\{\tau^{-2}\prod_{j=2}^{n}\frac{x\tau^{n-1}-z_j}{z_1\tau^2 - z_j} - \frac{x - z_1\tau^{n-1}}{x - z_1\tau^{n+1}}\prod_{j=2}^{n}\frac{x\tau^{-n+1}-z_j}{z_1\tau^2 - z_j}\right\}.
$$

Hence we get

$$
\begin{aligned}
&\tilde{\Delta}(z_1\tau^{n-1},x_2,\cdots,x_{n-2}|z_1,\cdots,z_n)\\
=\;&\prod_{j=2}^{n}\frac{z_j - z_1\tau^{2n-2}}{z_j - z_1\tau^2}\prod_{\mu=2}^{n-2}\frac{x_\mu - z_1\tau^{n-1}}{x_\mu - z_1\tau^{n+1}}\Delta(z_1\tau^{n+1},x_2,\cdots,x_{n-2}|z_1\tau^{2n},\cdots,z_n).
\end{aligned}
\tag{4.9}
$$

From Lemma 4.2 and (4.9), we obtain

$$
\tilde{\Delta}(z_1\tau^{n-1},x_2,\cdots,x_{n-2}|z_1,\cdots,z_n) = \prod_{j=2}^{n}\frac{z_j - z_1\tau^{2n-2}}{z_j - z_1\tau^2}\Delta(z_1\tau^{n-1},x_2,\cdots,x_{n-2}|z_1,\cdots,z_n). \tag{4.10}
$$

The polynomials $\tilde{\Delta}$ and Δ have the same factor $\prod_{1\le\mu\le\nu\le n-2}(x_\mu - x_\nu)$. Furthermore, the degrees of $\tilde{\Delta}$ and Δ with respect to x_μ are at most $(n-2)$. Thus we get

$$\begin{aligned} & \tilde{\Delta}(x_1, x_2, \cdots, x_{n-2}|z_1, \cdots, z_n) - \prod_{j=2}^{n} \frac{z_j - z_1\tau^{2n-2}}{z_j - z_1\tau^2} \Delta(z_1\tau^{n-1}, x_2, \cdots, x_{n-2}|z_1, \cdots, z_n) \\ = \; & c(z_1, \cdots, z_n) \prod_{1\le\mu<\nu\le n-2} (x_\mu - x_\nu) \prod_{\mu=1}^{n-2} (x_\mu - z_1\tau^{n-1}), \end{aligned} \tag{4.11}$$

where $c(z_1, \cdots, z_n)$ is homogeneous rational function of $(z_1, \cdots, z_n)$ whose total degree is 0. From Lemma 4.1 c has zeros at $z_j = z_1\tau^{2n-2}$, and may have poles at only $z_j = z_1\tau^2$. Thus c must have the form

$$c(z_1, \cdots, z_n) = s \prod_{j=2}^{n} \frac{z_j - z_1\tau^{2n-2}}{z_j - z_1\tau^2}, \qquad s \in \mathbb{C}.$$

By comparing both sides of (4.11) at $z_1 = 0$, we obtain $s = 0$. □

Therefore, by taking into account the symmetry (4.2), Theorem 3.1 for $m = 1$ was proved.

5 General case

In this section we prove Theorem 3.1 for general case. Let $I_l = \{(\gamma_1, \cdots, \gamma_l) | n \ge \gamma_1 > \cdots \gamma_l \ge 1\}$ for $1 \le l \le n-1$. Fix nonnegative integers $m(\gamma_1, \cdots, \gamma_l)$ for $1 \le l \le n-1$ and set

$$N_k = \sum_{l=1}^{n-1} \sum_{\substack{(\gamma_1, \cdots, \gamma_l) \in I_l \\ \gamma_j \ne k}} m(\gamma_1, \cdots, \gamma_l).$$

In what follows we often use the abbreviation $\gamma = (\gamma_1, \cdots, \gamma_l)$. Define the polynomial with respect to variables x and $\zeta^{(\gamma)} = (\zeta_1^{(\gamma)}, \cdots, \zeta_{m(\gamma)}^{(\gamma)})$ by

$$\begin{aligned} & A_\lambda(x|\zeta^{(n)}| \cdots |\zeta^{(1)}|\zeta^{(nn-1)}| \cdots |\zeta^{(21)}| \cdots |\zeta^{(\gamma_1 \cdots \gamma_l)}| \cdots |\zeta^{(n-1 \cdots 21)}) \\ = \; & \sum_{k=1}^{n} \prod_{l=1}^{n-1} \prod_{\substack{(\gamma)\in I_l, \\ \gamma_i = k}} \prod_{j=1}^{m(\gamma)} (x - \zeta_j^{(\gamma)} \tau^{2k-n-3+2i}) \\ \times \; & \frac{1}{x} f_\lambda^{(N_k)}(x\tau^{n+1-2k} | \bigcup_{l=1}^{n-1} \bigcup_{\substack{(\gamma)\in I_l, \\ \gamma_i > k > \gamma_{i+1}}} \bigcup_{j=1}^{m(\gamma)} \zeta_j^{(\gamma)} \tau^{2i-1}). \end{aligned} \tag{5.1}$$

For $n=3$ it reads as

$$\begin{aligned}
& xA_\lambda(x|\zeta^{(3)}|\zeta^{(2)}|\zeta^{(1)}|\zeta^{(32)}|\zeta^{(31)}|\zeta^{(21)}) \\
= & \prod_{j=1}^{m(3)}(x-\zeta_j^{(3)}\tau^2)\prod_{j=1}^{m(3,2)}(x-\zeta_j^{(32)}\tau^2)\prod_{j=1}^{m(3,1)}(x-\zeta_j^{(31)}\tau^2)f_\lambda^{(N_3)}(x\tau^{-2}|\zeta^{(2)}\tau^{-1},\zeta^{(1)}\tau^{-1},\zeta^{(21)}\tau^{-1}) \\
+ & \prod_{j=1}^{m(2)}(x-\zeta_j^{(2)})\prod_{j=1}^{m(3,2)}(x-\zeta_j^{(32)}\tau^2)\prod_{j=1}^{m(2,1)}(x-\zeta_j^{(21)})f_\lambda^{(N_2)}(x|\zeta^{(3)}\tau,\zeta^{(1)}\tau^{-1},\zeta^{(31)}\tau) \\
+ & \prod_{j=1}^{m(1)}(x-\zeta_j^{(1)}\tau^{-2})\prod_{j=1}^{m(3,1)}(x-\zeta_j^{(31)})\prod_{j=1}^{m(2,1)}(x-\zeta_j^{(21)})f_\lambda^{(N_1)}(x\tau^2|\zeta^{(3)}\tau,\zeta^{(2)}\tau,\zeta^{(32)}\tau^3).
\end{aligned}$$

It follows from the recursion relation

$$f_\lambda^{(N+1)}(y|z_1,\cdots,z_N,a)=f_\lambda^{(N)}(y|z_1,\cdots,z_N)-af_{\lambda-1}^{(N)}(y|z_1,\cdots,z_N),$$

that the polynomial $A_\lambda^{(m)}$ satisfies

$$\begin{aligned}
& A_\lambda^{(m)}(x|z^{(n)}|\cdots|z^{(1)})|_{\bigcup_{l=1}^{n-1}\bigcup_{j=1}^{l}z^{(\gamma_j)}=\zeta^{(\gamma_1\cdots\gamma_l)}\tau^{2j-2}} \\
= & \sum_{\rho=0}^{L}(-1)^\rho\sigma_\rho(\bigcup_{l=2}^{n-1}\bigcup_{\gamma\in I_l}\bigcup_{j=1}^{l-1}\zeta_j^{(\gamma)}\tau^{2j-1})A_{\lambda-\rho}(x|\zeta^{(n)}|\cdots|\zeta^{(1)}|\cdots|\zeta^{(n-1\cdots21)}),
\end{aligned}\tag{5.2}$$

where

$$\sum_{l=1}^{n-1}\sum_{\substack{(\gamma_1,\cdots,\gamma_l)\in I_l\\ \gamma_j=k}}m(\gamma_1,\cdots,\gamma_l)=m,\qquad (k=1,2,\cdots,n),$$

and

$$L=\sum_{l=2}^{n-1}(l-1)\sum_{\gamma\in I_l}m(\gamma).$$

For $n=3$ it reads as

$$\begin{aligned}
& A_\lambda^{(m)}(x|z^{(3)}|z^{(2)}|z^{(1)})|_{\bigcup_{l=1}^{2}\bigcup_{j=1}^{l}z^{(\gamma_j)}=\zeta^{(\gamma_1\cdots\gamma_l)}\tau^{2j-2}} \\
= & \sum_{\rho=0}^{L}(-1)^\rho\sigma_\rho(\zeta^{(32)}\tau,\zeta^{(31)}\tau,\zeta^{(21)}\tau)A_{\lambda-\rho}(x|\zeta^{(3)}\tau,\zeta^{(2)}\tau,\zeta^{(1)}\tau,\zeta^{(32)}\tau,\zeta^{(31)}\tau,\zeta^{(21)}\tau).
\end{aligned}$$

Define the polynomial for positive integer α

$$\begin{aligned}
& h_\alpha^{(m|L)}(x|\zeta^{(n)}|\cdots|\zeta^{(1)}|\zeta^{(nn-1)}|\cdots|\zeta^{(\delta_1\cdots\delta_l)}|\cdots|\zeta^{(n-1\cdots21)}) \\
= & x^{\alpha-2}\left\{\tau^{n(\alpha-1)}\prod_{l=1}^{n-1}\prod_{\gamma\in I_l}\prod_{j=1}^{m(\gamma)}(x\tau^{n-l(\gamma)}-\zeta_j^{(\gamma)}\tau^{l(\gamma)-1})\right. \\
- & \left.\tau^{-n(\alpha-1)}\prod_{l=1}^{n-1}\prod_{\gamma\in I_l}\prod_{j=1}^{m(\gamma)}(x\tau^{-n+l}-\zeta_j^{(\gamma)}\tau^{l(\gamma)-1})\right\}.
\end{aligned}$$

One of important observations is as follows:

Proposition 5.1 *For $m(\gamma_1,\gamma_2,\cdots,\gamma_l)$ such that $\sum_{l=1}^{n-1}\sum_{(\gamma_1,\cdots,\gamma_l)\in I_l,\gamma_j=k} m(\gamma_1,\cdots,\gamma_l)=m, (k=1,2,\cdots,n)$, the polynomial $\Delta^{(m)}$ satisfies the following relation.*

$$\begin{aligned}
&\Delta^{(m)}(x_1,\cdots,x_M|z^{(n)}|z^{(n-1)}|\cdots|z^{(1)})|_{\bigcup_{l=1}^{n-1}\bigcup_{j=1}^{l} z^{(\gamma_j)}=\zeta^{(\gamma_1\cdots\gamma_l)}\tau^{2j-2}}\\
=&\sum_{1\le\mu_1<\cdots<\mu_L\le M}(-1)^{\sum_{i=1}^{L}M-L+i+\mu_i}\det\left(h_{\alpha-1}^{(m)}(x_{\mu_i}|\zeta^{(n)}|\cdots|\zeta^{(1)}|\cdots|\zeta^{(n-1\cdots21)}\right)_{1\le i,\alpha\le L} \qquad (5.3)\\
\times&\ \Delta(\overset{x_\mu}{\hat{x}}\ |\zeta^{(n)}|\cdots|\zeta^{(1)}|\cdots|\zeta^{(n-1\cdots21)}).
\end{aligned}$$

Here

$$\begin{aligned}
&\Delta(y_1,\cdots,y_{M-L}|\zeta^{(n)}|\cdots|\zeta^{(1)}|\cdots|\zeta^{(n-1\cdots21)})\\
=&\ \det\left(A_\lambda(y_\mu|\zeta^{(n)}|\cdots|\zeta^{(1)}|\zeta^{(nn-1)}|\cdots|\zeta^{(21)}|\cdots|\zeta^{(n-1\cdots1)}\right)_{1\le\lambda,\mu\le M-L},
\end{aligned} \qquad (5.4)$$

and

$$L=\sum_{l=2}^{n-1}(l-1)\sum_{\gamma\in I_l}m(\gamma),\quad M=(n-1)m-1.$$

Proof This follows from the induction with respect to L. When $L=0$ (5.3) is obvious. Assume (5.3) for L. After another restriction, say $\zeta^{(1)}_{m(1)}=\zeta^{(2)}_{m(2)}\tau^2$, using (3.2) we have

$$\begin{aligned}
&A_{M-L}(x|\zeta^{(n)}|\cdots|\zeta^{(1)}|\zeta^{(nn-1)}|\cdots|\zeta^{(21)}|\cdots|\zeta^{(n-1\cdots1)})|_{\zeta^{(1)}_{m(1)}=\zeta^{(2)}_{m(2)}\tau^2}\\
=&\ h_1^{(m|L+1)}(x|\zeta^{(n)}|\cdots|\zeta^{(1)}|\zeta^{(nn-1)}|\cdots|\zeta^{(\delta_1\cdots\delta_l)}|\cdots|\zeta^{(n-1\cdots21)}).
\end{aligned}$$

Furthermore, we have

$$\begin{aligned}
&h_\alpha^{(m|L)}(x|\zeta^{(n)}|\cdots|\zeta^{(1)}|\cdots|\zeta^{(n-1\cdots1)})|_{\zeta^{(1)}_{m(1)}=\zeta^{(2)}_{m(2)}\tau^2}\\
=&\ h_{\alpha+1}^{(m|L+1)}(x|\zeta^{(n)}|\cdots|\zeta^{(1)}|\cdots|\zeta^{(n-1\cdots1)})-\zeta^{(2)}_{m(2)}\tau h_\alpha^{(m|L+1)}(x|\zeta^{(n)}|\cdots|\zeta^{(1)}|\cdots|\zeta^{(n-1\cdots1)}).
\end{aligned}$$

For other types of restrictions we get similar results. Thus using (5.2) and performing elementary transformation we have (5.3) for $L+1$. □

Thanks to the recursion relation for $f_\lambda^{(N)}$

$$f_\lambda^{(N+n-1)}(y|z_1,\cdots,z_N,a\tau,a\tau^3,\cdots,a\tau^{2n-3})=\sum_{\rho=0}^{n-1}(-1)^\rho\sigma_\rho(a\tau,a\tau^3,\cdots,a\tau^{2n-3})f_{\lambda-\rho}^{(N)}(y|z_1,\cdots,z_N),$$

the polynomial $A_\lambda^{(m)}$ satisfies the following recursion relation:

$$\begin{aligned}
&A_\lambda^{(m)}(x|z^{(n)}|\cdots|z^{(1)})|_{z_m^{(k)}=a\tau^{2n-2k}\quad(k=1,\cdots,n)}=(x-a\tau^{n-1})\\
\times&\ \sum_{\rho=0}^{n-1}(-1)^\rho\sigma_\rho(a\tau,a\tau^3,\cdots,a\tau^{2n-3})A_{\lambda-\rho}^{(m-1)}(x|z'^{(n)}|\cdots|z'^{(1)}).
\end{aligned} \qquad (5.5)$$

Note that for $1 \leq \alpha \leq n-1$

$$A^{(m-1)}_{M_{m-1}+\alpha}(z'^{(n)}|\cdots|z'^{(1)}) = h^{(m-1|0)}_{\alpha}(x|z'). \tag{5.6}$$

By combining (5.5) and (5.6) and using the same argument in the proof for Proposition 5.1 , one can show

Proposition 5.2 *The determinant $\Delta^{(m)}$ obeys the following recursion relation.*

$$\begin{aligned}
& \Delta^{(m)}(x_1,\cdots,x_M|z^{(n)}|\cdots|z^{(1)})\big|_{z^{(k)}_m=a\tau^{2n-2k},(k=1,2,\cdots,n)} \\
= & \prod_{\mu=1}^{M}(x_\mu - a\tau^{n-1}) \sum_{1\leq\mu_1<\cdots<\mu_{n-1}\leq M}(-1)^{\sum_{i=1}^{n} mn-m-n+i+\mu_i} \\
\times & \det\left(h^{(m-1|0)}_{\alpha}(x_{\mu_i}|z')\right)_{1\leq i,\alpha\leq n-1}\Delta^{(m-1)}(x\setminus\{x_\mu\}|z'),
\end{aligned}$$

where we use the abbreviation $x_\mu = (x_{\mu_1},\cdots,x_{\mu_{n-1}})$.

The following are two key theorems:

Theorem 5.3 *Let $P(m)$ be Proposition 3.3 for $m \geq 1$. Then for $m > 1$, $P(m)$ under the restriction $z^{(n)}_m = a,\cdots,z^{(2)}_m = a\tau^{2n-4}, z^{(1)}_{m-1} = a\tau^{2n-2}$ holds if $P(m-1)$ holds.*

Proof Using the relation

$$\varphi^{(m)}(j_1,\cdots,j_{n-1})\big|_{z^{(k)}_m=a\tau^{2n-2k},(k=2\cdots n),z^{(1)}_{m-1}=a\tau^{2n-2}} = (-\tau)^{n-1}\frac{a-z_0\tau^2}{a\tau^{2n-2}-z_0\tau^2}\varphi^{(m-1)}(j_1,\cdots,j_{n-1}),$$

the LHS of (3.6) can be expressed in a recursive way. Let us turn to the RHS. Note that

$$\begin{aligned}
& \tilde{h}^{(m|l)}_1(x|z'^{(n)}|\cdots|z'^{(n-l+1)}|z^{(n-l)}|\cdots|z^{(1)}|\cdots|\zeta^{(n\cdots n-l+1)}=a) \\
= & \frac{z_0\tau^{2n-2}-a\tau^{2l-4}}{z_0-a\tau^{2n-2}}\frac{1}{x}\left\{(x\tau^{n-l}-a\tau^{l-1})\prod_{\substack{j=0\\ j\neq km(k=1,\cdots,l)}}(x\tau^{n-1}-z_j)\right. \\
- & \left.(x\tau^{-n+l}-a\tau^{l-1})\prod_{\substack{j=0\\ j\neq km(k=1,\cdots,l)}}(x\tau^{-n+1}-z_j)\right\}.
\end{aligned} \tag{5.7}$$

From Proposition 5.2 and (5.7) we obtain

$$\begin{aligned}
& \tilde{\Delta}^{(m)}(x|z^{(n)}|\cdots|z'^{(1)},z_0)\big|_{z^{(k)}_m=a\tau^{2n-2k},(k=2\cdots n),z^{(1)}_{m-1}=a\tau^{2n-2}} \\
= & \prod_{l=2}^{n}\frac{z_0\tau^{2n-2}-a\tau^{2l-4}}{z_0-a\tau^{2l-4}}\prod_{\mu=1}^{M}(x_\mu-a\tau^{n-1})\sum_{1\leq\mu_1<\cdots<\mu_{n-1}\leq M}(-1)^{\sum_{i=1}^{n} mn-m-n+i+\mu_i} \\
\times & \det\left(h^{(m-1)}_{\alpha}(x_{\mu_i}|z'',z_0)\right)_{1\leq i,\alpha\leq n-1}\tilde{\Delta}^{(m-1)}(x\setminus\{x_\mu\}|z'',z_0),
\end{aligned}$$

where we use the abbreviation $z'' = (z'^{(n)}|\cdots|z'^{(2)}|z_1^{(1)},\cdots,z_{m-2}^{(1)})$. Thus the claim is verified. □

Theorem 5.4 *Let $P(m)$ be Proposition 3.3 for $m \geq 1$ Then $P(m)$ under the restriction $z_m^{(n)} = z_0\tau^2,\cdots,z_m^{(2)} = z_0\tau^{2n-2}$ holds.*

Proof Under the restriction in consideration there is only one nonzero term in the LHS of (3.6), and the only nonzero coefficient $\varphi^{(m)}$ is

$$\varphi^{(m)}(m,2m,\cdots,(n-1)m) = (-\tau)^{m(1-n)}\prod_{j=1}^{m-1}\frac{z_j'^{(1)} - z_0\tau^{2n}}{z_j'^{(1)} - z_0\tau^2}. \tag{5.8}$$

Since in the RHS there exists a zero and a pole under the restriction, we have to set $z_m^{(n)} = a,\cdots,z_m^{(2)} = a\tau^{2n-4}$ and reduce, then we have to set $a = z_0\tau^2$ to evaluate the RHS:

$$\begin{aligned}
&\tilde{\Delta}^{(m)}(x|z^{(n)}|\cdots|z^{(2)}|z'^{(1)},z_0)|_{z_m^{(k)}=z_0\tau^{2n+2-2k},(k=2,\cdots,n)}\\
=\;&(-\tau^{(n-1)})^{(n-2)}\prod_{k=2}^{n}\prod_{j=1}^{m-1}\frac{z_j^{(k)} - z_0\tau^{2n-2}}{z_j^{(k)} - z_0\tau^2}\prod_{k=2}^{n}\frac{z_j^{(1)} - z_0\tau^{2n}}{z_j^{(1)} - z_0\tau^2}\\
\times\;&\prod_{\mu=1}^{M}(x_\mu - z_0\tau^{n-1})\sum_{1\leq\mu_1<\cdots<\mu_{n-1}\leq M}(-1)^{\sum_{i=1}^{n}mn-m-n+i+\mu_i}\\
\times\;&\det(h_\alpha^{(m-1)}(x_{\mu_i}|z'))_{1\leq i,\alpha\leq n-1}\Delta^{(m-1)}(x\setminus\{x_\mu\}|z').
\end{aligned} \tag{5.9}$$

Therefore this proposition follows from (5.8) and (5.9). □

We now wish to show Proposition 3.3. First of all, note that LHS of Proposition 3.3 has no singularity at $z'^{(1)} = z_0\tau^{2n-2}$. Thus let us show the following proposition.

Proposition 5.5

$$\tilde{\Delta}^{(m)}(x_1,\cdots,x_M|z^{(n)}|z^{(n-1)}|\cdots|z'^{(1)},z_0)|_{z_j^{(k)}=z_0\tau^{2n-2}} = 0,\quad (k=2,\cdots,n)$$

Proof. By the same argument as in Proposition 5.2, we have an equation which can be obtained from (5.4) by replacing $A \to \tilde{A}$ and $\Delta \to \tilde{\Delta}$. Since the last row obtained in this way vanishes, the claim of this Proposition is verified. □

Using a general m analogue of (4.8), we can show

Proposition 5.6

$$\begin{aligned}
&\lim_{z_m^{(n)}\to z_0\tau^2}\prod_{k=2}^{n}\prod_{j=1}^{m}\frac{z_j^{(k)}-z_0\tau^2}{z_j^{(k)}-z_0\tau^{2n-2}}\tilde{\Delta}(x|z^{(n)}|\cdots|z^{(2)}|z'^{(1)},z_0)\\
=&\sum_{\nu=1}^{M}\prod_{\substack{\mu=1\\ \mu\neq\nu}}^{M}\frac{x_\mu-z_0\tau^{n-1}}{x_\mu-z_0\tau^{n+1}}g_1(x_\nu|z'^{(n)}|z^{(n-1)}|\cdots|z'^{(1)}|z_0)\\
\times&\ \Delta(x_1,\cdots,z_0\tau^{n+1}\overset{\nu}{\hat{}}\cdots x_M|z'^{(n)}|z^{(n-1)}|\cdots|z'^{(1)}|z_0),
\end{aligned}$$

where

$$g_1(x|z'^{(n)}|z^{(n-1)}|\cdots|z'^{(1)}|z_0)=\frac{z_0\tau^{n+1}}{x}\left\{\tau^{-2}\prod_{\substack{j=1\\ j\neq m}}^{N}\frac{x\tau^{n-1}-z_j}{z_0\tau^2-z_j}-\frac{x-z_0\tau^{n-1}}{x-z_0\tau^{n+1}}\prod_{\substack{j=1\\ j\neq m}}^{N}\frac{x\tau^{-n+1}-z_j}{z_0\tau^2-z_j}\right\}.$$

Now we are in a position to prove Proposition 3.3.

Proof of Proposition 3.3 From Proposition 5.5 both sides of (3.6) are homogeneous rational functions of degree $Mm+M(M-1)/2$ with simple poles located at $z'^{(1)}=z_0\tau^2$. Taking into account that the antisymmetry of x_μ's the equation (3.6) holds if both sides of (3.6) coincides at $(n-1)m^2$ points $z^{(n)}=z_0\tau^2$ and $z^{(a)}=z^{(b)}\tau^2(a<b)$. From Proposition 5.1 both sides of (3.6) after L times restriction can be expressed as follows:

$$\begin{aligned}
\mathrm{LHS}|_{\mathrm{restrictions}} &= \textstyle\sum_{1\le\mu_1<\cdots<\mu_L\le M}\det(h^{(m)}_{\alpha-1}(x_{\mu_i}|\zeta))_{1\le\alpha,i\le L}S_L(x\setminus\{x_\mu\}),\\
\mathrm{RHS}|_{\mathrm{restrictions}} &= \textstyle\sum_{1\le\mu_1<\cdots<\mu_L\le M}\det(h^{(m)}_{\alpha-1}(x_{\mu_i}|\zeta))_{1\le\alpha,i\le L}S_R(x\setminus\{x_\mu\}).
\end{aligned}$$

Here $S_L(y_1,\cdots,y_{M-L})$ and $S_R(y_1,\cdots,y_{M-L})$ are skew symmetric with respect to $(y_1,\cdots,y_{M-L})$. Note that the degrees of $S_L(y_1,\cdots)$ and $S_R(y_1,\cdots)$ with respect to y_1 are $M-L+m-1$. Furthermore, $\det(h^{(m)}_{\alpha-1}(y_1,y_2,\cdots,y_L|\zeta))_{1\le\alpha,j\le L}=cy_1^{M+m-1}y_2^{M+m-2}\cdots y_L^{M+m-L}+\cdots$. By comparing the coefficients of $\mathrm{LHS}|_{\mathrm{restrictions}}$ and $\mathrm{RHS}|_{\mathrm{restrictions}}$ with respect to $x_1^{M+m-1}x_2^{M+m-2}\cdots x_L^{M+m-L}$, we conclude that $\mathrm{LHS}|_{\mathrm{restriction}}=\mathrm{RHS}|_{\mathrm{restriction}}$ if and only if

$$S_L(x_1,\cdots,x_{M-L})=S_R(x_1,\cdots,x_{M-L}). \tag{5.10}$$

Consequently, taking into account that the antisymmetry of x_μ's, we have to only examine $\mathrm{LHS}|_{\mathrm{restriction}}=\mathrm{RHS}|_{\mathrm{restriction}}$ at $(n-1)m^2-Lm$ planes. After repeating this procedure the equation (3.6) reduces to Theorem 5.4 and 5.4. For simplicity and transparency, we first demonstrate how this induction scheme does work when $n=3$.

For $3\ge a>b\ge 1$, let $m(a,b)$ stand for the number of restrictions $z^{(b)}=z^{(a)}\tau^2$ and let $0\le l\le 1$ be the number of restriction $z^{(3)}=z_0\tau^2$. We denote such restriction by $(l,m(2,1)+m(3,2))$ and

introduce a lexicographical order by $(1,m) > (1,m-1) > \cdots > (1,0) > (0,m) > \cdots > (0,0)$. The restriction (l,k) is larger than (l',k') if $(l,k) > (l',k')$. We wish to show that $(l, m(2,1)+m(3,2))$ reduces to larger restrictions and $(1,m)$ reduces to Theorem 5.3 and 5.4.

1st step. Set $(l, m(2,1)+m(3,2)) = (1,m)$. In this case we need $m^2 - m(1+m(3,1))$ planes on which (5.10) holds for $L = m+1+m(3,1)$. We have m restrictions which reduce to Theorem 5.4, and $m(m-1) - m(2,1)m(3,2)$ restrictions which reduce to Theorem 5.3, and $(m-1-m(3,1)-m(3,2))(m-1-m(3,1)-m(2,1))$ restrictions which reduce to previous two restrictions. Hence we actually have $m^2 - m(1+m(3,1)) + (m(3,1)+1)^2$ planes.

2nd step. Set $(l, m(2,1)+m(3,2)) = (1,k)$, where $k < m$. In this case we need $2m^2 - m(1+k+m(3,1))$ planes on which (5.10) holds. We have m restrictions which reduce to Theorem 5.4, $k(m-1) - m(3,2)m(2,1)$ restrictions which reduce to Theorem 5.3, and $(m-k)(2m-k-2)$ restrictions which reduce to larger restrictions. We also have $(m-1-m(3,1)-m(3,2))(m-1-m(3,1)-m(2,1))$ additional restrictions at $z^{(1)} = z^{(3)}\tau^2$ which reduce to previous three restrictions. Hence we have $2m^2 - m(1+k+m(3,1)) + (k-m+1+m(3,1)/2)^2 + m(3,1)(3m(3,1)+4)/4$ planes.

3rd step. We can show in a similar way as in the 2nd step that this induction scheme does work for $l = 0$.

Therefore the equation (3.6) are proved for $n = 3$.

Next we show (3.6) for general n.

Let l denote the maximal length of restrictions such that $z^{(n+1-l)} = z^{(n+2-l)}\tau^2 = \cdots = z^{(n)}\tau^{2l-2} = z_0\tau^{2l}$. Let $r_j (1 \le j \le n-2)$ denote the number of restrictions of type $z^{(k)} = z^{(k+1)}\tau^2 = \cdots = z^{(k+j)}\tau^{2j}$, where $1 \le k \le n-j$. Set $L = \sum_{j=1}^{n-2} j r_j$. We introduce the lexicographical order in $\{(l, L, r_{n-2}, \cdots, r_1) \mid 0 \le L + l \le (n-2)m + 1\}$ by

$$(n-2, (n-2)(m-1)+1, m-1, 0, \cdots, 0, 1) > (n-2, (n-2)(m-1)+1, m-2, 1, 0, \cdots, 0, 2)$$
$$> \cdots > (0,1,0,\cdots,0,1) > (0,0,0,0,\cdots,0).$$

At the stage of induction of type $(l, L, r_{n-2}, \cdots, r_1)$, we need more than or equal to $(n-1)m^2 - m(l+L)$ restriction planes.

Remark 1. In the expression of (5.1) let us call z which belongs to one of $\zeta^{(i)}$ *free* z. Then we have at least $m-2$ free z's.

Remark 2. There are restrictions which are not counted by $r_j (j = 1, \cdots, n-2)$; e.g., restrictions counted by $m(a,b)$, where $a - b > 1$. Let us call such a restriction a *bad restriction*. Even if we need less than or equal to m bad restrictions to obtain $(n-1)m^2 - m(l+L)$ restriction planes for $(l, L, r_{n-2}, \cdots, r_1)$, our induction scheme does work. Because after a bad restriction, we have

still as the same number of not bad restrictions as that for $(l, L, r_{n-2}, \cdots, r_1)$, while we need only $(n-1)m^2 - m(l+L) - m$ restriction planes.

1st step. Set $(l, L, r_{n-2}, \cdots, r_1) = (n-2, (n-2)(m-1)+1, m-1, 0, \cdots, 0, 1)$. Then we need $m^2 - m$ planes on which (5.10) holds. Let us first consider $m(n, \cdots, 2) = m-1, m(2,1) = 1, m(1) = m-2$, and the other $m(\gamma) = 0$. We have m planes which reduce to Theorem 5.4, and $(m-1)^2$ planes which reduce to Theorem 5.3. Thus we have actually $m^2 - m + 1$ planes. Next consider $m(n, \cdots, 2) = m-2$ and $m(n-1, \cdots, 1) = 1$. In this case we have m planes which reduce to Theorem 5.4, and $(m-1)^2 - (m-2)$ planes which reduce to Theorem 5.3. We also have $m-2$ bad restrictions. Hence from Remark 2 this case reduces to Theorems 5.3 and 5.4. In general when we replace $(m(n, \cdots, 2), m(n-1, \cdots, 1)) = (m-p, p-1)$ by $(m(n, \cdots, 2), m(n-1, \cdots, 1)) = (m-1-p, p)$, we need $m - 2p$ additional bad restrictions. Thus every $(l, L, r_{n-2}, \cdots, r_1)$ reduces to Theorems 5.3 and 5.4. In a similar way, one can show that for other restrictions of type $(l, L, r'_{n-2}, \cdots, r'_1)$ we have $m^2 - m + 1$ planes while we need $m^2 - m$ planes.

2nd step. Set $(l, L) = (n-2, (n-2)(m-1))$. Such a restriction can be obtained by dividing one of chains

$$z^{(k)} = z^{(k+1)}\tau^2 = \cdots = z^{(k+j)}\tau^{2j}, \tag{5.11}$$

of a restriction of type $(l, L+1)$ into two pieces like

$$z^{(k)} = \cdots = z^{(k+i)}\tau^{2i}, \quad \text{and} \quad z^{(k+i+1)}\tau^{2(i+1)} = \cdots = z^{(k+j)}\tau^{2j}, \tag{5.12}$$

where $0 \le i \le j-1$.

Note that there are two kinds of restriction planes for (l, L), *old ones* and *new ones* : old ones are restriction planes which decrease the degree of (3.6) even for (5.11); and new ones are those which does not decrease the degree of (3.6) for (5.11) but does decrease it for (5.12).

For example, the following restriction of type (l, L)

$$\begin{aligned}
&z_1^{(3)} = z_1^{(4)}\tau^2 = \cdots = z_1^{(n)}\tau^{2(n-3)} = z_0\tau^{2(n-2)},\\
&z_i^{(2)} = z_i^{(3)}\tau^2 = \cdots = z_i^{(n)}\tau^{2(n-2)}, \quad 2 \le i \le m-1,\\
&z_m^{(2)} = z_m^{(3)}\tau^2 = \cdots = z_m^{(n-1)}\tau^{2(n-3)},\\
&z_1^{(1)} = z_1^{(2)}\tau^2,
\end{aligned} \tag{5.13}$$

can be obtained from the following restriction of type $(l, L+1)$

$$\begin{aligned}
&z_1^{(3)} = z_1^{(4)}\tau^2 = \cdots = z_1^{(n)}\tau^{2(n-3)} = z_0\tau^{2(n-2)},\\
&z_i^{(2)} = z_i^{(3)}\tau^2 = \cdots = z_i^{(n)}\tau^{2(n-2)}, \quad 2 \le i \le m,\\
&z_1^{(1)} = z_1^{(2)}\tau^2,
\end{aligned} \tag{5.14}$$

by resetting the relation between $z_m^{(n-1)}$ and $z_m^{(n)}$. For (5.14) we have m^2-m+1 restriction planes

$$\begin{aligned} z_j^{(2)} &= z_1^{(3)}\tau^2, \quad 1 \le j \le m, \\ z_k^{(1)} &= z_1^{(2)}\tau^2, \quad 2 \le i, \le m, 1 \le k \le m-1, \end{aligned} \tag{5.15}$$

which reduce to Theorems 5.3 and 5.4. For (5.13) m^2-m+1 restriction planes (5.15) are old ones. On the other hand, restriction planes $z_m^{(n-1)} = z_m^{(n)}\tau^2$, $z_1^{(2)} = z_m^{(n)}\tau^2$ and $z_k^{(1)} = z_m^{(n)}\tau^2 (1 \le k \le m-1)$ are new ones.

It is evident that the number of restriction planes for $(l, L+1)$ and that of old restriction planes for (l, L) coincide. Thus in order to show this case, we need $m^2-(m^2-m+1) = m-1$ new restriction planes.

Let us consider the case $i=0$ in (5.12). Now there are p free z's belonging to $\zeta^{(k)}$, and there are $m-1-p$ free z's belonging to $\zeta^{(i)}$, where $i \neq k$, and $1 \le p \le m-1$. Thus we have at least $p+p(m-1-p) \ge m-1$ new restriction planes. One can show the other case $1 \le i \le j-1$ similarly. Hence this reduces to larger restrictions.

3rd step. Suppose that we have enough number of restriction planes to show (5.10) for a restriction of type $(n-2, L, r_{n-2}, \cdots, r_1)$, where $L \le (n-2)(m-1)$. If we replace L by $L-1$, we need m new restriction planes. On the other hand, there are now at least m free z's. Thus, by a parallel argument given in *2nd step* this case reduces to larger restriction.

4th step. Suppose that we have enough number of restriction planes to show (5.10) for the restriction of type $(l, L, r_{n-2}, \cdots, r_1)$, where $L \le (n-2)(m-1)$. If we replace l by $l-1$, we need m new restriction planes. Now we have at least m free z's. Thus, from the same argument given in *2nd step*, this case reduces to larger restrictions. As for the restriction obtained by replacing L by $L-1$, the argument again perfectly parallels the one given in *2nd step*.

Therefore, after repeating this procedure, we reach

$$(l, L, r_{n-2}, \cdots, r_1) = (0, \cdots, 0),$$

and this is what we wish to prove. □

Next we notice that Proposition 3.3 implies (1.7) for $\varepsilon_1 = i$. For that purpose let us seek the residue formula for $\overline{G}$.

Let

$$G(z^{(n)}|\cdots|z^{(1)}) = \oint_C dx_1 \cdots \oint_C dx_M \overline{G}(x|z^{(n)}|\cdots|z^{(1)})\Psi(x|z),$$

and

$$\overline{G}_i(z^{(n)}|\cdots|z^{(1)}) = G^{\overbrace{n\cdots n}^{m}\cdots\hat{i}\cdots\hat{i}\cdots\overbrace{1\cdots 1}^{m}\overbrace{i\cdots i}^{m}}(z^{(n)}|\cdots|z^{(1)}).$$

Then we have

$$\begin{aligned}\overline{G}_i^{(m)}(x|z) &= (-\tau)^{i-1} \sum_{1\le\mu_1<\cdots<\mu_{n-1}\le M} (-1)^{\sum_{i=1}^n mn-m-n+i+\mu_i} \\ &\times \det\left(h_\alpha^{(m-1)}(x_{\mu_i}|z')\right)_{1\le i,\alpha\le n-1} \overline{G}_i^{(m-1)}(\overset{\mu}{\hat{x}}|z).\end{aligned} \tag{5.16}$$

Therefore (3.4) for $i \neq 1$ reduces to Proposition 3.3.

Thus we have proved the Theorem 3.1.

6 Conclusions and discussions

In this paper we present an integral formula for quantum KZ equation of level 0 associated with the vector representation of $U_q(\widehat{\mathfrak{sl}_n})$. This work is a generalization of our previous paper [1]. Smirnov obtained the formula for form factors of the $SU(n)$ chiral Gross-Neveu model in Appendix A in his book [3]. It gives a rational scaling limit of the present work (i.e., $q = e^\epsilon,\ z = e^{-\epsilon n\frac{\beta}{\pi i}}, \epsilon \to 0$). Smirnov [3] has studied form factors of integrable massive theories. Equations (1.1) and (1.2) are two of the axioms proposed by him that form factors obey. Let $G(z)$ be a form factor of a certain operator. Then a ϑ function in the integral kernel should be determined. It is interesting and important to solve this problem.

Finally, we discuss the related works on the integral formulas for the quantum KZ equations. In [5] an integral formula for correlation functions of the XXZ model was obtained on the basis of the bosonization of the level 1 highest weight representations of $U_q(\widehat{\mathfrak{sl}_2})$. This scheme gives only one particular solution though those correlations satisfy the quantum KZ equation of arbitrarily level. A formula for higher spin analog of the XXZ model in terms of Jackson-type integral was given in [6], by using the level k bosonization of $U_q(\widehat{\mathfrak{sl}_2})$.

In [7, 8] solutions by Jackson-type integrals are obtained. Their formulae are in principle valid for general level, as opposed to our integral formula restricted to level 0. On the other hand, the problem of choosing the cycles for Jackson-type integrals, which accommodates the freedom of the solutions, is not well studied.

Tarasov and Varchenko [9] improved the Jackson-type integral formula for $U_q(\mathfrak{gl}_n)$ such that the solutions automatically satisfy the R-matrix symmetry. The number of Jackson-type integrals is $n(n-1)m/2$ for $N = nm$, whereas our formula is written by $(n-1)m-1$ fold integral. For $n = 2$ two numbers are different by only 1, however, the difference gets greater as n increases.

Acknowledgments

The authors would like to thank M. Jimbo, T. Miwa and F. A. Smirnov for a number of discussion. Y.-H.Q. is partly supported by the Grant-in-Aid for Scientific Research from the Ministry of Education, Science and Culture, Japan, No. 04-2297; and by Soryushi Shogakukai.

References

[1] Jimbo, M., Kojima, T., Miwa, T. and Quano, Y.-H., Smirnov's integrals and quantum Knizhnik–Zamolodchikov equation of level 0, J. Phys. **A27** (1994) 3267–3283.

[2] Frenkel, I. B. and Reshetikhin, N. Yu., Quantum affine algebras and holonomic difference equations, Comm. Math. Phys. **146** (1992) 1-60.

[3] Smirnov, F. A., *Form Factors in Completely Integrable Models of Quantum Field Theory*, World Scientific, Singapore, 1992.

[4] Smirnov, F. A., Dynamical symmetries of massive integrable models 1,2: Int. J. Mod. Phys. **7A, Suppl. 1B** (1992) 813-837, 839-858.

[5] Jimbo, M., Miki, K., Miwa, T. and Nakayashiki, A., Correlation functions of the XXZ model for $\Delta < -1$, Phys. Lett. **168A** (1992) 256-263.

[6] Kato, A., Quano, Y.-H. and Shiraishi, J., Free boson representation for q-vertex operators and their correlation functions, Comm. Math. Phys. **157** (1993) 119-137.

[7] Reshetikhin, N. Yu., Jackson-type integrals, Bethe vectors, and solution to a difference analog of Knizhnik–Zamolodchikov system, Lett. Math. Phys. **26** (1992) 153-165.

[8] Matsuo, A., Quantum algebra structure of certain Jackson integrals, Comm. Math. Phys. **157** (1993) 479-498.

[9] Tarasov, V. and Varchenko, A., Jackson integral representations for solutions of the quantized Knizhnik–Zamolodchikov equation, RIMS preprint, RIMS-949.